## Bryn Mawr Medical Specialists Endoscopy Center Clenpiq Colonoscopy Preparation-<u>Afternoon Procedure</u>

Appointment Date: \_\_\_\_\_ Arrival Time\_\_\_\_\_

7 Days Prior	5 Days Prior	3 Days Prior	2 Days Prior	1 Day Prior	Procedure Day
Arrange a ride. You will be sedated for this procedure & must have an escort, over the age of 18, to drive you home. You are not permitted to take a cab or Uber ride home, unless you are accompanied by a responsible adult (family or friend).  Please make necessary arrangements or speak with our office regarding ride concerns.  You are not permitted to drive, work, exercise, operate equipment or drink alcohol, for the remainder of the day, after your procedure.  If you are diabetic, please inform your Primary Care Physician or Endocrinologist about your scheduled procedure and discuss your diabetic medications. Adjustments to your medications must be made.	Stop taking the following medications: Vitamins, Herbal Supplements, Ibuprofen (Advil, Motrin) or Aleve.  If you take Aspirin because you have risk factors for coronary artery disease or stroke: DO NOT stop Aspirin  If you take other blood thinners, such as Coumadin or Plavix, carefully follow the instructions, provided at the time of scheduling, regarding need to hold them. You may be asked to contact your cardiologist for special instructions to hold blood thinners.  Check with your insurance company for the following: -Benefits -Deductible -Co-Pay for outpatient surgery (Your final co-pay responsibility will be determined after your claim is submitted.)	*Last Chance to cancel your procedure without a \$100 cancellation fee: physician office (610) 525-9570.  Review upcoming dietary needs for the next 2 days.  Purchase Clenpiq with prescription.  Disregard instructions on the box & follow these instructions.  Stop eating nuts, seeds and berries.	Drink at least 8 glasses of water throughout the day.  Begin a "Low residue Diet" today (see handout).  Please confirm your ride.	Start "Clear Liquid Diet" today (see handout).  No Red or purple liquids  At 5pm, drink one bottle of Clenpiq. Follow with (5) 8 oz drinks of clear liquids within the next 4 hours.  Please continue to drink additional clear liquids throughout the day and evening.  If you vomit the preparation, please call the Physician on call at (610) 526-0881.	Take heart and blood pressure medications.  At 6am drink a bottle of Clenpiq. Follow with (5) 8oz drinks of clear liquids.  The bowel prep must be completed 4 hours prior to your arrival time. Then nothing by mouth.  Be sure to bring your completed paperwork, current insurance card and photo ID with you.  Your driver must be present in the endoscopy suite, at the time of discharge.  Bryn Mawr Medical Specialists Endoscopy Center is located in Suite 330 of the Bryn Mawr Medical Arts Pavilion at 825 Old Lancaster Road, Bryn Mawr-NOT at Bryn Mawr Hospital.

Rev. 12/2018