Bryn Mawr Medical Specialists Endoscopy Center Miralax Colonoscopy Preparation-<u>Afternoon Procedure</u>

7 Days Prior	5 Days Prior	3 Days Prior	2 Days Prior	1 Day Prior	Procedure Day
Arrange a ride. You will be sedated for this procedure & must have an escort, over the age of 18, to drive you home. You are not permitted to take a cab or Uber ride home, unless you are accompanied by a responsible adult (family or friend). Please make necessary arrangements or speak with our office regarding ride concerns. You are not permitted to drive, work, exercise, operate equipment or drink alcohol, for the remainder of the day, after your procedure. If you are diabetic, please inform your Primary Care Physician or Endocrinologist about your scheduled procedure and discuss your diabetic medications. Adjustments to your medications must be made.	Stop taking the following medications: Vitamins, Herbal Supplements, Ibuprofen (Advil, Motrin) or Aleve. If you take Aspirin as requested by a cardiologist: DO NOT stop Aspirin If you take Aspirin for general health and not because of cardiac reasons: DO stop Aspirin If you take blood thinners, such as Coumadin or Plavix, carefully follow the instructions, provided at the time of scheduling, regarding need to hold them. You may be asked to contact your cardiologist for special instructions to hold blood thinners. Check with your insurance company for the following: -Benefits -Deductible -Co-Pay for outpatient surgery Co-Pays and deductibles are paid the day of your procedure.	*Last Chance to cancel your procedure without a \$100 cancellation fee: physician office (610) 525-9570. Review upcoming dietary needs for the next 2 days. Purchase the following: Miralax: 8.3 oz. bottle (238 gram, 14 day) Sold over the counter. Dulcolax laxative 5 mg. tablets (Bisacodylgeneric) You'll need 4 tablets. Sold over the counter. 64 ounces of Gatorade or low sugar drink: Propel Fitness Water, Gatorade 2, or Powerade Zero. *No red or purple clear liquids. Stop eating nuts, seeds and berries.	Drink at least 8 glasses of water throughout the day. Begin a "Low residue Diet" today (see handout). Remember to fill out your online health history from the link sent to your email address. If you do not have access to email or have trouble accessing it, a nurse will call you 1 day prior to your appointment and take your health history over the phone. Please allow time for this phone call.	Start "Clear Liquid Diet" today (see handout). No solid food today. After your clear liquid "breakfast", mix the 8.3 oz bottle of Miralax with 64 ounces of Propel or Gatorade. Chill in Refrigerator. At 4 pm take (2) Dulcolax (bisacodyl) tablets At 6pm, begin drinking the Miralax mixture. Drink an 8oz glass every 15-30 minutes until ½ the mixture is finished. At 8 pm take (2) Dulcolax (bisacodyl) tablets Please continue to drink additional clear liquids throughout the day and evening. If you vomit or have difficulty with the bowel preparation, please call the Physician on call: (610) 526-0881.	Take heart and blood pressure medications. At 6am begin drinking the remainder of the Miralax mixture. Drink an 8 oz glass every 15-30 minutes until finished The bowel prep must be completed 4 hours prior to your arrival time. Then nothing by mouth. No Cannabis use of any form after midnight. Be sure to bring your current insurance card and photo ID with you. Your driver must be present in the endoscopy suite at the time of discharge. Bryn Mawr Medical Specialists Endoscopy Center is located in Suite 330 of the Bryn Mawr Medical Arts Pavilion at 825 Old Lancaster Road, Bryn Mawr-NOT at Bryn Mawr Hospital.