## Bryn Mawr Medical Specialists Endoscopy Center SUFLAVE Colonoscopy Preparation-<u>Morning Procedure</u>

Appointment Date: \_\_\_\_\_

Arrival Time\_\_\_\_\_

7 Days Prior	5 Days Prior	3 Days Prior	2 Days Prior	1 Day Prior	Procedure Day
Arrange a ride. You will be sedated for this procedure & <u>must</u> have an escort, over the age of 18, to drive you home. You are <u>not</u> permitted to take a cab or Uber ride home, unless you are accompanied by a responsible adult (family or friend). Please make necessary arrangements or speak with our office regarding ride concerns. You are not permitted to drive, work, exercise, operate equipment or drink alcohol, for the remainder of the day, after your procedure. If you are diabetic, please inform your Primary Care Physician or Endocrinologist about your scheduled procedure and discuss your diabetic medications. Adjustments to your medications must be made.	Stop taking the following medications:Vitamins, HerbalSupplements, Ibuprofen (Advil, Motrin) or Aleve.If you take Aspirin as requested by a cardiologist: DO NOT stop AspirinIf you take Aspirin for general health and not because of cardiac reasons: DO stop AspirinIf you take blood thinners, such as Coumadin or Plavix, carefully follow the instructions, provided at the time of scheduling, regarding need to hold them. You may be asked to contact your cardiologist for special instructions to hold blood thinners.Check with your insurance company for the following: -Benefits -Deductible -Co-Pay for outpatient surgery Co-Pays and deductibles are paid the day of your procedure.	<ul> <li>*Last Chance to cancel your procedure without a \$100 cancellation fee: physician office (610) 525-9570.</li> <li>Review upcoming dietary needs for the next 2 days.</li> <li>Purchase <u>SUFLAVE</u> with prescription. If you do not have your <u>SUFLAVE</u> yet, please call the office to verify your. 610-525-9570</li> <li>*IMPORTANT Disregard instructions inside the <u>SUFLAVE</u> box &amp; follow <u>these</u> instructions.</li> <li>Stop eating nuts, seeds and berries.</li> </ul>	Drink at least 8 glasses of water throughout the day. Begin a "Low residue Diet" today (see handout). Remember to fill out your online health history from the link sent to your email address. If you do not have access to email or have trouble accessing it, a nurse will call you 1 day prior to your appointment and take your health history over the phone. Please allow time for this phone call.	Start "Clear Liquid Diet" today (see handout). No Red or purple liquids SUFLAVE DOSING INSTRUCTIONS: Add 1 flavor enhancing packet to each of the two bottles provided. Fill each bottle with lukewarm water up to the fill line. After capping the bottle, gently shake each bottle until all powder is mixed well. For best taste, refrigerate for at least one hour. Use within 24 hours. At <b>5pm</b> , take 1 bottle from the fridge and drink 8 ounces of solution every 15 minutes until the bottle is empty. Drink an additional 16 ounces of water before next dose. At <b>9pm</b> , take the second bottle from the fridge and drink as instructed in previous step. Drink an additional 16 ounces of water.	Stop drinking liquids at least 4 hours prior to your colonoscopy.NOTHING TO EAT UNTIL AFTER YOUR COLONOSCOPYTake your heart and blood pressure medications by 6am with a small sip of water.Be sure to bring your current insurance card and photo ID with you.Your driver must be present in the endoscopy suite at the time of discharge.Bryn Mawr Medical Specialists Endoscopy Center is located in Suite 330 of the Bryn Mawr Medical Arts Pavilion at 825 Old Lancaster Road, Bryn Mawr Hospital.If you vomit the preparation, please call the Physician on call at (610) 526-0881.